Fountaine, M. E. 1902. Butterfly hunting in Greece in 1900. *Ent. Record.* **14**: 12-35; 64-67.

Higgins, L. G. and Riley, N. D. 1983. *A Field Guide to the Butterflies of Britain and Europe* (5th Edition). Collins, London.

Hurtig, F. 1940. Nuovi contributi alla conoscenza della fauna delle Isole Italiane dell Egeo, XII. *Boll. Labour. Ent. Postici.* **3**: 221-246.

Koutsaftikis, A. 1974. Recent butterfly records from Greece. *Ent. Record.* **86**: 15-17.

Larsen, T. B. 1974. *Butterflies of Lebanon.* Publication of the National Council for Scientific Research (C.N.R.S.), Beirut, Lebanon.

Rebel, H. 1936. Zoologische Ergebnisse einer Dodekanesreise von O. Wettsbein 1935. Lepidoptera. *S.B.A.K.Wiss. Wein.* 145: 19-33.

Straubenzee, C. van. 1932. Three months collecting in Greece. *Entomologist* **65**: 154-159; 177-183.

Thomson, G. 1969. *Maniola (Epinephele) jurtina* (L.) (Lep.) Satyridae) and its forms. *Ent. Record* **51**: 51-58.

Turati, E. 1929. Ricerche faunistiche nell "Isole Italiane dell" Egeo. Lepidotteri. *Archo. Zool. Ital.* **13**: 178-186.

A BELATED RECORD OF BISELACHISTA TRAPEZIELLA STAINTON (LEP.: ELACHISTIDAE) FOR SCOTLAND. — Further to Dr. K. P. Bland's interesting not on this species (cf. *Entomologist's Rec. J. Var.* **98**: 27), I wish to record that in April 1973, I collected mines and larvae of *B. trapeziella* in Argyllshire in the town of Oban and nearby on Pulpit Hill, all in *Luzula sylvatica*, but unfortunately failed to breed any. However, I kept the mines and recently on submitting them to Dr. R. Knill-Jones, who knows the species well, he kindly confirmed the determination. — J. M. CHALMERS-HUNT.

CELAENA HAWORTHII CURT. (HAWORTH'S MINOR) AND EUPITHECIA TENUIATA HUBN. (SLENDER PUG) ON JERSEY 1984 — Single specimens of these two species were caught in the Rothamsted Insect Survey light trap on Jersey (Site No. 146) on 4-ix-84 and 20-vii-84 resp. and were determined by examination of the genitalia. So far as I am aware *C. haworthii* and *E. tenuiata* have not previously been recorded from the Channel Islands.

Thanks are extended to Mrs. R. Collier and Mr. G. Thomas for operating the Jersey trap and to Mr. R. Long of the Societe Jersiaise for his comments on these two species. ADRIAN M. RILEY, Entomology Department, Rothamsted Experimental Station, Harpenden, Herts.